

A New Species of Columbellid Gastropod from the Old World Tropics

William K. Emerson

Department of Invertebrates
American Museum of Natural History
New York, NY 10024-5192, USA

ABSTRACT

Cotonopsis monfilsi new species, is described from off Senegal, West Africa in 230-500 m. The generic assignment is made on the basis of shell, opercular, and radular characters. The radula is typically columbellid, whereas the shell morphology lacks some characteristics of the "Strombina" gastropods. Therefore, the placement in the *Strombina*-group genus *Cotonopsis* is tentative. *Metula dockeryi* new name is proposed to replace *Metula inflata* Dockery, 1984, not *Metula inflata* (Houbrick, 1984).

Key Words: Prosobranchia, Columbellidae, *Cotonopsis*, Buccinidae, *Metula*, west Africa, new species, new name.

INTRODUCTION

During the past five years, we have received from several shell dealers specimens of an apparently new species of columbellid gastropod for study and report. The taxonomic placement of these shells could not be made with certainty until the morphology of the radula was determined. A radula was eventually found and proved to be typically columbellid in form. As a result, this gastropod can be assigned provisionally to the *Strombina*-group genus *Cotonopsis*.

I take pleasure in describing this new species in honor of Paul Monfils, who was the first to call this perplexing species to my attention.

A review of the literature disclosed that a replacement name was required for a fossil species of *Metula*. A new name is provided for *Metula inflata* Dockery, 1984, not *Metula inflata* Houbrick, 1984.

ABBREVIATIONS

AMNH = American Museum of Natural History, New York
LACM = Los Angeles County Museum of Natural History, California.

SYSTEMATICS

Superfamily: Buccinoidea Rafinesque, 1815

Family Columbellidae Swainson, 1840


Genus *Cotonopsis* Olsson, 1942

Type species (by original designation): *Strombina (Cotonopsis) panacostariceus* Olsson, 1942, Pliocene of the Burica Peninsula, Costa Rica, Charco Azul Formation

Cotonopsis monfilsi new species
(figures 1-10)

Description: Shell solid, fusiform, whorls inflated without a sutural ramp; protoconch of 3½ smooth, convex whorls; teleoconch of 7½ convex whorls, sculptured with numerous fine spiral lirae, numbering about 18 on the penultimate whorl and about 32 on the body whorl. Some spiral lines bifid; lirae not consistently evenly spaced. Thickened varix formed on body whorl above the rim of the aperture. Aperture ovate, inner lip terminally thin, but with a sublirial ridge with 10 lirae teeth; columella with 4-5 weak folds; siphonal canal open, short, narrow. Fasciole indistinct. Periostracum brownish tan. Background color brownish buff-tan with broken whitish bands below the suture and with axial blotches interrupting the tanish coloration. Aperture white. Operculum thin, corneous, lenticular with a terminal nucleus at edge. Radular ribbon (figs. 9, 10) rachiglossate, typically columbellid (cf. Radwin, 1977, fig. 22b). Rachidian tooth a narrow, nearly rectangular plate. Lateral tooth shaft-like with barbed base, tip of shaft with two hooked cusps.

Material examined: Holotype (AMNH 232519) and 3 paratypes (AMNH 232091) from the type locality; 5 Paratypes (AMNH 232092) dredged in 300-350 m, on muddy bottom, off St. Louis, Senegal, November, 1987, ex-Northeast Natural History Imports; 3 Paratypes (AMNH 232152) dredged in 450-500 m, on muddy-sand bottom, off St. Louis, Senegal, ex-Northeast Natural History Imports; 6 referred specimens (5 specimens, AMNH




Figures 1-8. *Cotonopsis monfilsii* n. sp. 1-3, holotype (AMNH 232519). 4-6, paratype (AMNH 232091); figs. 1-6 $\times 1.5$. Figs. 7, 8, paratype (AMNH 232092), spire enlarged to show detail of early whorls; figs. 7, 8, $5.0\times$.

232176; 1 specimen, LACM 146956) dredged on silty-sand bottom, 230-260 m, off Casamance Province, Senegal ($12^{\circ}50'N, 15^{\circ}0'W$), ex-Northeast Natural History Imports; 2 referred specimens (AMNH 255056) trawled in deep water from off the Dakar harbor entrance, Senegal ($14^{\circ}40'N, 17^{\circ}26'W$), ex-Mal de Mer Enterprises. See Table 1 for shell measurements.

Type locality: dredged in 300 m off St Louis, Senegal, West Africa, ($16^{\circ}02'N, 16^{\circ}30'W$) by Marcel Pin, 1987, ex-Abbey Specimen Shells.

Distribution: Known only from off Senegal, West Africa in 230 to 500 meters.

Remarks: The shells of the present specimens resemble certain "*Strombina*"-group gastropods (such as *Cotonopsis*), but they are also in some respects reminiscent of certain buccinid gastropods (eg. *Pisania*, *Metula*, etc.). Before the nature of the radula was known, I asked several colleagues for their opinions based on the examination of photographs or the study of the specimens. Some of them referred the specimens to the buccinid genus *Pisania* Bivona-Bernardi, 1832, in the subfamily Pisaniinae Gray, 1857:13, formerly credited to Tryon, 1881:98 (see Cernohorsky, 1971:138; 1975:192, Beu and Maxwell, 1987:56; and Bouchet, 1988:149, for comments on the subfamilial status of Pisaniinae in Buccinidae, and



Figures 9, 10. *Cotonopsis monfilsii* n. sp. 9, holotype (AMNH 232519), radula, about one-third of the ribbon. 10, detail of the rachidian teeth; fig. 9, scale bar = 100 μ m, fig. 10, scale bar = 200 μ m, courtesy of J. H. McLean

its rejection without an explanation at the subfamilial level by Ponder and Warén, 1988:305). The resemblance of the new species to Recent species of *Charitodoron* (Mitridae) from South Africa was also suggested (see Lozouet, 1991:206, figs. 26–30). Other colleagues believed the Senegalese specimens to be an undescribed columbellid species referable to *Cotonopsis* (*sensu stricto*).

With the recovery of a radula from a dried body of the new species, the placement of the new species in the family Columbellidae can be confirmed. The radular characters are typically columbellid in form (figs. 9, 10). As in *Cotonopsis argentca* (Houbrick, 1983:352, fig. 2), the radular dentition consists of rachidian teeth that are thin, narrow and form a rectangular plate. A wide space separates the rachidian teeth from the lateral teeth, which are composed of a shaft with an enlarged base and are tipped with two hooked cusps. The shells of the new species possess the thickened varices of *Strombina*-group gastropods, but the apertural dentition of the inner lip differs in having evenly spaced lirate denticles of equal size. These are comparable to those of the buccinid genus *Bartschia* Rehder (1943:199, pl. 20, fig. 17; Olsson and Bayer, 1972:924, fig. 14). Furthermore, *Bartschia significans*, the type species, has strongly cancellate sculpture, whereas the new species possesses fine spiral lirae. The radular characters of *Bartschia* are unknown, but on shell features, Beu and Maxwell (1987:62) believe *Bartschia*, together with *Metula* H. and A. Adams (1853: 84; Emerson, 1986:27), to be closely related to the buccinid genus *Colubraria* Schumacher, 1817. Some of the western Atlantic species of *Metula* [e.g. the West Indian *M. (Agassitula) agassizi* Clench and Aguayo, 1941:179, pl. 14, fig. 4; Olsson and Bayer, 1972:917, fig. 11 and the east African *M. (Kanamarua) rehderi* (Kilburn, 1977: 193, fig. 21)] superficially resemble the new species. Bouchet (1988:150, fig. 1) illustrated the typically buc-

Table 1. Shell dimensions and proportions of the specimens of *Cotonopsis monfilsii* n. sp. n = 20 specimens. Width includes terminal varix on body whorl. Measurements in mm.

Type specimens	Height	Width	Width-height
*AMNH 232519 (holotype)	40.1	14.9	0.37
AMNH 232091 (paratype 1)	41.7	15.2	0.36
AMNH 232091 (paratype 2)	37.2	13.7	0.37
AMNH 232091 (paratype 3)	37.1	13.6	0.37
AMNH 232092 (paratype 1)	41.1	14.4	0.35
AMNH 232092 (paratype 2)	37.9	13.4	0.35
AMNH 232092 (paratype 3)	37.1	13.8	0.35
AMNH 232092 (paratype 4)	37.5	13.6	0.36
AMNH 232092 (paratype 5)	36.2	12.8	0.35
AMNH 232152 (paratype 1)	40.4	14.6	0.36
AMNH 232152 (paratype 2)	36.9	12.8	0.35
AMNH 232152 (paratype 3)	34.5	12.6	0.37
Referred specimens			
AMNH 232176, a	39.7	14.5	0.37
AMNH 232176, b	36.3	13.8	0.38
AMNH 232176, c	36.1	13.9	0.39
AMNH 232176, d	35.0	14.0	0.40
AMNH 232176, e	34.9	13.7	0.40
LACM 146956	36.2	13.9	0.38
AMNH 255056, a	39.1	14.1	0.36
AMNH 255056, b	38.6	13.4	0.35
Mean	37.1	13.8	0.36
Range	34.5–	12.6–	0.35–
	41.7	15.2	0.40

* Radula extracted.

cinid radula of his new species, *Metula africana*, and concluded the radular morphology is significantly different from that of the buccinid genus *Pisania*.

The placement of the new species in the genus *Cotonopsis* based on shell characters is tentative. As the genus is defined by Jung (1989:158), the teleoconch whorls are smooth or axially sculptured. In the new species, the teleoconch sculpture is composed of fine spiral lines and no axial sculpture is present. *Cotonopsis* is represented in the tropical western Atlantic by one species (*C. argentca*). The remainder of the 15 species of *Cotonopsis* recognized by Jung (1989), 2 fossil and 12 living species, occur in the west American tropical zone. Previously, the *Strombina*-group gastropods were believed to be restricted to the New World tropics, dating from the Miocene to the present, with only 3 of the 35 living species occurring in the western Atlantic (Jung, 1989:4). Therefore, the presence of a species of *Cotonopsis* in west African waters may reflect a separate lineage of these columbellids that may eventually be determined to require recognition as a new genus-group taxon.

REPLACEMENT NAME FOR A HOMONYM

During my review of the literature for this paper, I noted a homonymic species-group name in *Metula* (Buccinidae). *Metula (Metula) inflata* Dockery in MacNeil and

Dockery (1984:331, pl. 52, fig. 3; published in "November, 1984", *teste* D. T. Dockery III, in *litt.*, Dec. 9, 1991) is a junior secondary homonym of *Acamptochetus* [= *Metula*] *inflatus* Houbriek (1984:421, fig. 1; published July 6, 1984). I here rename *Metula inflata* Dockery, 1984 (not *Metula inflata* [Houbriek, 1984]), as *Metula dockeryi* Emerson, NEW NAME. Dockery's taxon is a Paleogene fossil from Mississippi and Houbriek's taxon is a Recent Philippine species.

Acamptochetus Cossmann (1901:123; type species by original designation: *Murex mitraeformis* Brocchi, 1814) is a junior subjective synonym of *Metula* H. and A. Adams (1853:84; type species by subsequent designation Kobelt, 1876: *Buccinum clathratum* Adams and Reeve, 1850, [not Kiener, 1834, nor Anton, 1839] = *Metula amosi* Vanatta, 1913:22), as noted by Emerson (1986:27) and Beu and Maxwell (1987:62) and accepted by Bouchet (1988:149).

ACKNOWLEDGEMENTS

In addition to Paul Monfils of Northeast Natural History Imports, Providence, RI, I am grateful to Robert Foster of Abbey Specimen Shells of Santa Barbara, CA and Robert and Dorothy Janowsky of Mal De Mer Enterprises, West Hempstead, NY for providing specimens for study. Silvard P. Kool of the Museum of Comparative Zoology, Harvard University, kindly lent specimens of *Metula* for comparisons. The following colleagues cooperatively offered comments on the taxonomic placement of the new species before the nature of the radula was known: Walter O. Cernohorsky, M. G. Harasewych, Richard S. Houbriek, Peter Jung, R. N. Kilburn, Jorgen Knudsen, Paula M. Mikkelsen, Gary Rosenberg, and Graham Saunders. James H. McLean generously mounted the radula and provided the photographs for the radula. I am indebted to my AMNH colleagues, Walter E. Sage, III for technical assistance, Andrew S. Modell for photographic services and Stephanie Crooms for word-processing the manuscript.

LITERATURE CITED

Adams, H. and A. 1853 [-1852]. The Genera of recent Mollusca, arranged according to their organization, London 1(13):65-96, pls. 9-12 (June, 1853).

Anton, H. E. 1839. Verzeichniss der Conchylien welche sich in der Sammlung von Hermann Eduard Anton befinden. Halle, xvi + 110 p.

Beu, A. G. and P. A. Maxwell. 1987. A revision of the fossil and living gastropods, related to *Plesiotriton* Fischer, 1884 (Family Cancellariidae, subfamily Plesiotritoninae n. subfam.) with an Appendix: Genera of Buccinidae Pisaninae related to *Colubraria* Schumacher, 1817. New Zealand Geological Survey Paleontological Bulletin 54 1-140.

Bivona-Bernardi, A. 1832. Caratteri d'un nuovo genere di conchiglie della famiglia delle columellarie del Signor De Lamarck. Effemeridi Scientifiche e Letterarie per la Sicilia 2:8-13, 1 pl.

Bouchet, P. 1988. Two new species of *Metula* (Gastropoda

Buccinidae) with a description of the radula of the genus. The Nautilus 102(4):149-153.

Cernohorsky, W. O. 1971. Indo-Pacific Pisaninae (Mollusca: Gastropoda) and related buccinid genera. Records of the Auckland Institute and Museum 8:137-167.

Cernohorsky, W. O. 1975. Supplementary notes on the taxonomy of buccinid species of the subfamily Pisaninae (Mollusca: Gastropoda). Records of the Auckland Institute and Museum 12:175-211.

Clench, W. J. and C. G. Agnayo. 1941. Notes and descriptions of new deep-water Mollusca obtained by the Harvard-Havana Expedition off the coast of Cuba. Memorias de la Sociedad Cubana de Historia Natural, Havana 15(2):177-180.

Cossmann, M. 1901. Essais de Paléconchologie Comparée, pt 4, 293 p.

Emerson, W. K. 1986. On the type species of *Metula* H. and A. Adams, 1853: *Buccinum clathratum* A. Adams and Reeve, 1850 (Gastropoda: Buccinidae). The Nautilus 100(1): 27-30.

Gray, J. E. 1857. Guide to the systematic distribution of Mollusca in the British Museum. Part 1, London, 230 p.

Houbriek, R. S. 1983. A new *Strombina* species (Gastropoda: Prosobranchia) from the tropical western Atlantic. Proceedings of the Biological Society of Washington 96(3): 349-354.

Houbriek, R. S. 1984. A new "*Metula*" species from the Indo-West Pacific (Prosobranchia Buccinidae). Proceedings of the Biological Society of Washington 97(2):420-424 (published July 6, 1984).

Jung, P. 1989. Revision of the *Strombina*-Group (Gastropoda: Columbellidae), fossil and living. Distribution, biostratigraphy, systematics. Mémoires Suisses de Paléontologie 111: 1-298.

Kiener, L. C. 1834 [-1841]. Spécies général et iconographie des coquilles vivantes. . . Paris, 9:1-112.

Kilburn, R. N. 1977. Taxonomic studies on the marine Mollusca of southern Africa and Mozambique. Part 1. Annals of the Natal Museum 23(1):173-214.

Kobelt, W. 1876 [-1881]. Illustriertes Conchylienbuch. Nuremberg, 1:1-143.

Lozouet, P. 1991. Mollusca Gastropoda: *Eumitra* récentes de la région néo-calédonienne et *Charitodoron* fossiles de l'Oligocène supérieur d'Aquitaine (Mitridae). In: Crosnier, A. and P. Bouchet (eds.). Résultats des Campagnes MUSORSTOM, vol 7, no. 8. Mémoires de la Muséum National d'Histoire Naturelle, Ser. A, Zoologie 150:205-222.

MacNeil, F. S., and D. T. Dockery, III. 1984. Lower Oligocene Gastropoda, Scaphopoda, and Cephalopoda of the Vicksburg Group in Mississippi. Mississippi Department of Natural Resources, Bureau of Geology, Bulletin 24:1-415 (published in November, 1984).

Olsson, A. A. 1942. Tertiary and Quaternary fossils from Burica Peninsula of Panama and Costa Rica. Bulletin of American Paleontology 27(106):153-258.

Olsson, A. A. and F. M. Bayer. 1972. American metulas (Gastropoda: Buccinidae). Bulletin of Marine Science (University Miami, FL) 22(4):900-925.

Ponder, W. F. and A. Warén. 1988. In: Ponder, W. F., D. J. Eernisse, and J. H. Waterhouse, eds. Prosobranch Phylogeny. Appendix, Classification of the Caenogastropoda and Heterostropha—A list of the family-group names and higher taxa. Malacological Review, Supplement 4:288-326.

Radwin, G. E. 1977. The family Columbellidae in the western Atlantic. The Veliger 19(4):403-417.

Rehder, H. A. 1943. New marine mollusks from the Antillean region. Proceedings of the United States National Museum 93(3161):187-203, pls. 19, 20.

Tryon, G. W. [1880-] 1881. Manual of conchology; structural and systematic. Series 1, Tritonidae, Fusidae, Buccinidae,

3:1-310, Academy of Natural Sciences, Philadelphia [p. 98, 1881].

Vanatta, E. G. 1913. Descriptions of new species of marine shells. Proceedings of the Academy of Natural Sciences, Philadelphia 65(1):22-27, 3 text figs., pl. 2.